

# DEADLINE<sup>®</sup>

## DIFENACOU M PASTE BAIT

HIGHLY PALATABLE PASTE BAIT FOR PROFESSIONAL USE AGAINST MICE.  
FOR USE IN INDOOR AND OUTDOOR SITUATIONS.

300g e

### DIRECTIONS FOR USE:

Wear gloves and any other suitable personal protective equipment. Use only in the caulking gun supplied. Cut off the dome end of the tube, taking care not to damage the screw thread. Cut off the tapered plastic nozzle to achieve an opening of approximately 1cm in diameter. Screw the nozzle onto the tube before installing the caulking gun. Apply up to 4g of bait into both corners of the bait box. After application, deposit any 'run-on' by drawing the end of the nozzle against the bait container. Cover the nozzle when not in use and release the caulking gun pressure to prevent leakage. Place bait boxes where mice are active, in runs or near harbourages. As a guide, place boxes 5 metres apart for high infestations and 10 metres apart for low infestations. Boxes containing the bait should not be placed near hot pipes and ovens etc. Always protect baits well so as to protect from the prevailing weather and to prevent access by humans, domestic animals and wildlife. Inspect baiting points at appropriate intervals and replenish bait as necessary until no more bait is taken. For high infestations, revisit within the first 7 days and then fortnightly until the infestation is under control. Thereafter, revisit as necessary. For low infestations, revisit within the first 7-10 days. Thereafter, revisit as necessary.

**Notes:** Difenacou m Paste Bait should be used between 16°C and 25°C (between 60°F and 77°F) for optimum use - outside these temperatures the product may be too hard or too soft.

**PRECAUTIONS:** To avoid risks to man and the environment, comply with the instructions for use. Unless under the supervision of a pest control operator or other competent person, do not use anticoagulant rodenticides as permanent baits. Keep locked up and out of the reach of children. Store in a cool, dry place. Prevent access to bait by children, birds and non-target animals (particularly dogs, cats, pigs and poultry). Baits must be securely deposited in a way so as to minimise the risk of consumption by other animals or children. Where possible, secure baits so that they cannot be dragged away. When tamper resistant bait stations are used, they should be clearly marked to show that they contain rodenticides and that they should not be disturbed. When the product is being used in public areas, the areas treated must be marked during the treatment period and a notice explaining the risk of primary and secondary poisoning by the anticoagulant as well as indicating the first measures to be taken in case of poisoning must be made available alongside the baits. Wash hands and exposed skin before meals and after use. Do not place bait where food, feed or water could become contaminated. Time for the biocidal effect is 3-21 days depending on the amount of bait consumed. The time to control an infestation completely will depend on factors such as the size of the infestation and the potential for rodents to come in from the surrounding areas. Search for and remove dead rodents at frequent intervals during treatment, at least as often as when baits are checked and/or replenished.

**FIRST AID:** This product contains an anticoagulant compound. If ingested, symptoms may include nosebleed and bleeding gums. In severe cases, there may be bruising, haematomas and blood present in the faeces and urine. In case of accident or if you feel unwell, seek medical advice immediately and show the label where possible.

**Active Ingredient:** DIFENACOU M 0.005% w/w

**NOTE TO DOCTOR:** Difenacou m is an indirect anticoagulant. Phytomenadione, Vitamin K1, is antidotal. Determine prothrombin times not less than eighteen hours after consumption. If elevated, administer Vitamin K1 and continue until prothrombin time normalises. Continue determination of prothrombin times for two weeks after withdrawal of the antidote and resume treatment if elevation occurs in that time. For further information contact the National Poisons Information Service on 0844 892 011 (for use by medical professionals only).

**DISPOSAL:** All pesticides must be disposed of as hazardous waste. This product must be disposed of as hazardous waste. Empty containers may be disposed of as controlled waste in accordance with the appropriate regulations. Dispose of dead rodents in accordance with local requirements. Remove all baits and bait trays after treatment and dispose of them in accordance with local requirements.

**AUTHORISATION NUMBER: UK-2011-0148**

CLD41  
011112-004C  
Batch No. See pack

**Rentokil Initial Supplies**  
Liverpool L33 7SR  
Enquiries: 0151 548 5050  
products@rentokil.com  
www.rentokilproducts.com  
Emergency line: 01342 833022



The bitter taste of Bitrex<sup>®</sup> helps to prevent accidental ingestion of this product.



Deadline is a registered trademark of Rentokil Initial plc.